

submitted to Astrophysical Journal Letters

Star Formation at the Twilight of the Dark Ages: Which Stars Reionized the Universe?

Rachel S. Somerville & Mario Livio ¹

*Space Telescope Science Institute
3700 San Martin Dr., Baltimore MD 21218*

ABSTRACT

We calculate the global star formation rate density (SFRD) from $z \sim 30$ – 3 using a semi-analytic model incorporating the hierarchical assembly of dark matter halos, gas cooling via atomic hydrogen, star formation, supernova feedback, and suppression of gas collapse in small halos due to the presence of a photoionizing background. We compare the results with the predictions of simpler models based on the rate of dark matter halo growth and a fixed ratio of stellar-to-dark mass, and with observational constraints on the SFRD at $3 \lesssim z \lesssim 6$. We also estimate the star formation rate due to very massive, metal-free Pop III stars using a simple model based on the halo formation rate, calibrated against detailed hydrodynamic simulations of Pop III star formation. We find that the total production rate of hydrogen-ionizing photons during the probable epoch of reionization ($15 \lesssim z \lesssim 20$) is approximately equally divided between Pop II and Pop III stars, and that if reionization is late ($z_{\text{reion}} \lesssim 15$, close to the lower limit of the range allowed by the *WMAP* results), then Pop II stars alone may be able to reionize the Universe.

Subject headings: cosmology:theory — galaxies:evolution — intergalactic medium

1. Introduction

When and how the Universe became reionized is one of the fundamental questions in cosmology. There is currently a great deal of interest in this question due to recent observational and theoretical developments. The discovery of several $z \sim 6$ quasars in the

¹somerville, mlivio@stsci.edu

SDSS whose spectra are consistent with showing zero flux below Lyman- α (a ‘Gunn-Peterson’ trough) may indicate that the IGM was predominantly neutral at $z \gtrsim 6$ (Fan 2001; Becker et al. 2001). The ionization history of the Universe may also be constrained via observations of the cosmic microwave background (CMB). The recently released first-year results from the *WMAP* satellite constrain the reionization redshift to the range $z_{\text{reion}} = 17 \pm 5$ (Kogut et al. 2003).

What is the nature of the sources that produced the photons responsible for this transition from neutral to ionized? In Cold Dark Matter (CDM) models, halos large enough to form significant amounts of molecular hydrogen start appearing around redshift $z \sim 30$ (Tegmark et al. 1997). The stars that formed in these halos, out of pristine, metal-free and magnetic-field-free gas, with molecular hydrogen as the only coolant, were almost certainly quite different from the stars we see around us today. Numerical simulations suggest that these first stars were extremely massive, on the order of a few hundred solar masses (Abel et al. 2000, 2002; Bromm et al. 2002). This first generation of very massive, metal free stars is often referred to as ‘Pop III’. Because of their high temperatures and low metallicities, Pop III stars may produce up to ~ 20 times as many hydrogen-ionizing photons per solar mass as Pop II stars (Bromm et al. 2001). It is therefore natural to think that these stars may play an important role in early reionization. Ironically, less is known about the *second* generation of stars, precisely because the metals, magnetic fields, and photons produced by the first stars complicate the situation. The key processes that determine how efficiently Pop III stars can form and when the transition to Pop II occurs — destruction of H_2 by Lyman-Werner photons, catalysis of H_2 by X-rays, and the production, dispersal, and mixing of heavy elements — remain poorly understood (Machacek et al. 2001, 2003; Ricotti et al. 2001, 2002; Cen 2002; Yoshida et al. 2003).

Around $z \sim 30$ –20, halos that are large enough to cool by atomic processes start to collapse. Two processes are likely to be responsible for regulating star formation in these halos. If the Universe has already been reionized, the UV background will prevent gas from collapsing into halos with temperatures smaller than the effective Jeans mass, $\sim 2 \times 10^5$ K, corresponding to circular velocities $V_c \sim 30$ –50 km/s (e.g. Thoul & Weinberg 1996; Gnedin 2000). We hereafter refer to the suppression of star formation resulting from this effect as photoionization ‘squenching’. Supernovae and massive stars also regulate star formation by heating the ISM and driving winds that remove the gas from the galaxy (Dekel & Silk 1986; Mac Low & Ferrara 1999).

In this *Letter*, we present predictions for the cosmic star formation rate density (SFRD) due to both Pop II and Pop III stars from $z \sim 30$ –3. We compare our results with observations at the low redshift end ($z \sim 3$ –6). We use a semi-analytic model to explore the effects of photoionization squenching and supernova feedback on these predictions, and consider sev-

eral possible reionization histories (including a model with multiple reionizations). We then assess the relative contribution from Pop II and Pop III stars to the ionizing photon budget during the redshift range relevant to the epoch of reionization as constrained by *WMAP* and the *SDSS* quasar observations ($z \sim 6\text{--}30$). Throughout, we assume cosmological parameters consistent with the *WMAP* data (Spergel et al. 2003): matter density $\Omega_m = 0.3$, baryon density $\Omega_b = 0.044$, dark energy $\Omega_\Lambda = 0.70$, Hubble parameter $H_0 = 70$ km/s/Mpc, fluctuation amplitude $\sigma_8 = 0.9$, and a scale-free primordial power spectrum $n_s = 1$.

2. The halo-collapse model

In order for gas to be converted into stars, the first-order condition is that it must be inside a collapsed halo of sufficient mass to allow the gas to cool and become dense. Following this line of argument, we may model the global star formation rate density by assuming that it is proportional to the rate at which gas collapses into halos in a given mass range:

$$\dot{\rho}_* = e_* \rho_b \frac{dF_h}{dt}(M > M_{\text{crit}}), \quad (1)$$

where $\frac{dF_h}{dt}(M > M_{\text{crit}})$ is the time derivative of the fraction of the total mass in collapsed halos with masses greater than M_{crit} , obtained from the halo mass function $dn_h/dM(M, z)$ given by the Press-Schechter model (Press & Schechter 1974) or one of its variants, and ρ_b is the mean density of baryons. One may then assign the efficiency of conversion of gas into stars e_* and the critical mass M_{crit} for separate populations depending on the main coolant and the mode of star formation. For example, M_{crit} corresponds to a halo virial temperature of about 10^4 K for halos that cool via atomic processes, while $T_{\text{crit}} \simeq 100$ K for molecular cooling. Pop III stars are generally assumed to form in the lower temperature, H_2 -cooled halos, with a much lower efficiency than Pop II stars, which are associated with larger, H_I -cooled halos (for a more detailed discussion, see e.g. Loeb & Barkana 2001). We hereafter refer to this as the ‘halo-collapse’ model.

3. Merger Tree models

Variants on simple models like the one presented above have been used in many recent analytic studies of early star formation and reionization (e.g. Cen 2002; Wyithe & Loeb 2002, 2003; Venkatesan et al. 2003; Haiman & Holder 2003). However, there is a well-developed semi-empirical approach to modeling the physics of atomic cooling, Pop II star formation and chemical enrichment, and supernova feedback, within the framework of hierarchical

merging predicted by CDM models. The effect of photoionization squelching on Pop II star formation has also been included in some semi-analytic models (Kauffmann et al. 1993; Somerville 2002; Benson et al. 2002). This approach has been used in a large number of studies of galaxy formation at lower redshift $z \sim 0\text{--}5$ (e.g. Kauffmann et al. 1993; Cole et al. 1994; Kauffmann et al. 1998; Cole et al. 2000; Somerville & Primack 1999; Somerville et al. 2001). It is interesting to see how the results based on these more realistic recipes compare with the simple ‘halo-collapse’ model described above, and to study the relative importance of the various processes that are expected to regulate star formation at very high redshift.

Here we use the models developed in Somerville & Primack (1999, SP) and Somerville et al. (2001, SPF), with photoionization squelching added as described in Somerville (2002), using a recipe based on the numerical results of (Gnedin 2000). We follow halo merger histories down to halos with temperature $T_{\text{vir}} = 10^4$ K, where atomic cooling becomes possible. We shall refer to these models as the ‘merger tree’ models.

4. Results

4.1. The Cosmic Star Formation History

We show the predicted star formation rate density (SFRD) for Pop II and Pop III stars in Fig. 1. For Pop III stars, we have used the halo collapse model (Eqn. 1) with $e_*^{\text{III}} = 0.001$ and $M_{\text{crit}}^{\text{III}} = 1.0 \times 10^6 h^{-1} M_{\odot}$. Also shown in Fig. 1a are the results from detailed numerical hydrodynamic simulations of Pop III star formation in a cosmological volume by Yoshida et al. (2003), for an assumed characteristic Pop III mass of $100 M_{\odot}$. We see that with these parameter choices, our simple recipe reproduces the simulation results fairly well. Of course, we do not really expect Pop III star formation to continue to $z \sim 3$. However, in the absence of more detailed modeling of metal production and mixing, we do not know when the Pop III mode will ‘shut off’.

Fig. 1a also shows the prediction of the halo collapse model for Pop II stars, with $M_{\text{crit}}^{\text{II}}$ corresponding to a temperature² of 10^4 K and efficiencies of $e_*^{\text{II}} = 0.1$ and $e_*^{\text{II}} = 1.0$. The $e_*^{\text{II}} = 1$ halo collapse model represents a sort of upper limit for star formation in halos that cool via H_I , as it would require all new gas flowing into these halos to cool and form stars instantaneously. The predicted SFRD may be compared with observational estimates and limits on this quantity at ‘low’ redshift $3 \lesssim z \lesssim 6$ (see figure caption). The assumed efficiencies of $e_*^{\text{II}} = 0.1$ and $e_*^{\text{II}} = 1$ approximately bracket the range in observational

²See SP for the conversions between virial mass, velocity, and temperature as a function of redshift

estimates. We note in passing that the Lanzetta et al. high estimate exceeds even the extreme case of $e_*^{\text{II}} = 1$ at $z = 8\text{--}10$.

We also show the star formation history produced in the more realistic semi-analytic merger tree models described in Section 3. To study the sensitivity of our results to our recipes for star formation, photoionization squelching, and supernova feedback, we investigate different choices of ingredients and parameters. The merger tree models are summarized in Table 1. The reader only interested in the main result may skip directly to Section 4.2.

Models MT-1a and MT-1b do not include supernova feedback or photoionization squelching. Star formation is then regulated only by the rate at which gas can cool via atomic processes and collapse, and by the star formation timescale $\tau_* = m_{\text{cold}}/\dot{m}_*$. We consider two choices for this timescale, which we refer to as ‘constant efficiency’ and ‘accelerating’. In the constant efficiency (CE) recipe, τ_* is a constant (we take $\tau_* = 0.1$). In the ‘accelerating’ recipe, $\tau_* \propto t_{\text{dyn}}$, where $t_{\text{dyn}} \propto r_{\text{vir}}/v_{\text{vir}}$ is the dynamical time of the halo. This is similar to the scaling observed in nearby galaxies (Kennicutt 1983, 1998) and is commonly used as an empirical basis for star formation recipes in semi-analytic models and hydrodynamic simulations. Referring to Fig. 1a, we see that the ‘accelerating’ scaling leads to dramatically more star formation at high redshift compared with the constant efficiency scaling. We use this recipe in the rest of the models.

In models MT-2a–c, we include the effect of photoionization squelching (but no SN feedback), with different assumed reionization histories, motivated by the joint *WMAP* and *SDSS* constraints discussed above. We show in Fig. 1a a model in which hydrogen is reionized at $z_{\text{reion}} = 20$ or $z_{\text{reion}} = 15$ and remains reionized, and a model with a ‘double reionization’ based on the model of Cen (2002), in which the Universe is reionized at $z_{\text{reion}}^{(1)} = 15.5$, recombines by $z_{\text{rec}} = 13.5$, and experiences a second reionization at $z_{\text{reion}}^{(2)} = 6$. The ‘dips’ in the star formation history are due to the suppression of gas infall by the photoionizing background, and demonstrate that a substantial fraction of the total star formation in the Universe would be taking place in small mass halos in the absence any kind of feedback.

Models MT-3a–c, shown in Fig. 1b, all assume the Cen (2002) double-reionization history described above, and also include supernova feedback. Note that the ‘dips’ are now much less dramatic, because much of the gas has already been removed from the small halos that are affected by photoionization squelching. An uncertain aspect of implementing supernova feedback in semi-analytic models is the fate of the reheated gas. In model MT-3a, the gas reheated by SN is removed from the disk but retained in the halo (‘retention’ feedback), in MT-3b the reheated gas is removed from the disk *and* dark matter halo of all galaxies (‘ejection’ feedback), and in model MT-3c the gas is ejected from the halo only if the halo virial velocity is less than 100 km/s. This threshold for ejection of gas by super-winds is

motivated by theoretical arguments (Dekel & Silk 1986) and observations of nearby galaxies (Martin 1999). The global star formation rate changes by as much as a factor of six at $z \sim 3$ depending on these choices, but by less than a factor of two at $z \gtrsim 15$. Note that the ‘dip’ following the first reionization is considerably more pronounced in the model with ejection feedback, as gas which has been accreted before reionization and ejected is not allowed to re-collapse in small halos while the photoionizing background is ‘switched on’. We consider models MT-3a and MT-3c to be the most realistic of the models considered here. Models with similar ingredients and parameter values have been shown to reproduce the luminosity function of galaxies at $z \sim 0$ (SP) as well as of Lyman break galaxies at $z \sim 3$ (SPF).

4.2. Production of Ionizing Photons

In Fig. 2 we show the cumulative number of hydrogen-ionizing photons per hydrogen atom in the Universe, produced by Pop II stars in our fiducial models (MT-3a and MT-3c), and by our model for Pop III star formation. For Pop II, we use the results of Leitherer et al. (1999) for the number of $\lambda < 912 \text{ \AA}$ photons produced by low metallicity stars with a Salpeter IMF. For Pop III, we assume that each star produces $1.6 \times 10^{48} \text{ photons s}^{-1} M_{\odot}^{-1}$ for a lifetime of 3 million years (Bromm et al. 2001). We emphasize that we have shown the number of ionizing photons *produced*, without attempting to fold in the fraction of these photons that manage to escape from the galaxy or to propagate through the IGM. The ionization fraction x_e is expected to scale as this quantity times $f_{\text{esc}} f_{\text{ion}} / C_{\text{clump}}$ (e.g., Spergel et al. 2003), where f_{ion} is the number of ionizations per UV photon, f_{esc} is the fraction of ionizing photons that escape from the galaxy, and C_{clump} is the clumping factor, reflecting the clumpiness of the IGM. The recombination time at $z \sim 10 - 25$ is $\sim 0.2 - 0.7$ times the Hubble time (Cen 2002). Observational constraints on f_{esc} for both nearby and high redshift galaxies range from a few percent to ~ 50 percent (Leitherer et al. 1995; Hurwitz et al. 1997; Steidel et al. 2001). Clumping factors at high redshift $z \gtrsim 10$ are expected to be on the order of $C_{\text{clump}} = 2 - 10$ (Cen 2002). Therefore the combination $f_{\text{esc}} f_{\text{ion}} / C_{\text{clump}}$ is plausibly of order $\sim 0.1 - 0.2$ in the relevant redshift range.

In Fig. 2, we see that the total number of ionizing photons produced by Pop II stars in our fiducial model overtakes the contribution from Pop III stars at around $z \sim 16 - 12$. If reionization occurred near the lower redshift end of the *WMAP* range (which seems easier to reconcile with our model), then our results suggest that Pop II stars contributed at least half of the ionizing photons. If reionization occurred as late as $z_{\text{reion}} \sim 12$, then Pop II stars (which are known observationally to exist) may even have been able to reionize the Universe on their own. One should also keep in mind that given the non-negligible rate of star formation (both Pop III and Pop II) even at $z \gtrsim 20$, we might expect there to have

been sufficient metal pollution to have halted the formation of very massive Pop III stars at quite a high redshift. However, the redshift of this transition is highly uncertain, and is almost certainly not sharp, but will occur at different epochs in different environments.

5. Discussion

We have shown predictions of the global star formation history from the time when the first stars began to form $z \sim 30$ until the epoch of the most distant observed galaxies $z \sim 3$ –6. We summarize our main conclusions as follows:

- There are two main competing effects determining the Pop II star formation efficiency (SFE) at high redshift. (i) The star formation activity shifts toward smaller halos at earlier times, leading to decreased effective SFE as supernova feedback and photoionization squelching reduce the available cold gas supply in these halos. (ii) If the SFE is higher when the gas density is larger, as suggested by observations in nearby galaxies, then this is partially counteracted by the increased efficiency expected due to the higher characteristic gas densities.
- Our results suggest that the background of hydrogen-ionizing photons during reionization at $z \sim 15$ –20 is roughly equally divided between Pop II and Pop III stars. While Pop III stars produce a larger number of ionizing photons per baryon, this is compensated by the lower SFRD of Pop III stars in the relevant redshift range $z \sim 15$ –20.
- If z_{reion} is pushed to the lower limit of the range allowed by the *WMAP* results, the background produced by Pop II stars alone may be sufficient to reionize the universe, thus removing the only compelling *observational* argument for the existence of Pop III stars.

Using similar semi-analytic recipes for star formation applied within N-body simulations with radiative transfer, Ciardi et al. (2003) also concluded that Pop II stars alone can produce reionization histories that are within the *WMAP* constraints.

There are several uncertain factors that could shift our results by factors of a few in different directions. The characteristic masses of the Pop III stars could be a factor of 2–3 higher than we have assumed here, resulting in a corresponding scaling in the star formation and ionizing photon production rate. However, we may have underestimated the number of ionizing photons produced by Pop II stars at early times, as they may have lower metallicities than the Leitherer et al. (1999) models used here, and may have a somewhat top-heavy IMF (Larson 1998). This could lead to a factor of ~ 2 –3 increase in the number of ionizing photons

predicted by the Pop II stars (Tumlinson et al. 2003; Schaerer 2003). It is clear that more detailed modeling of chemical evolution is needed to better constrain these uncertainties, as well as the transition redshift between massive Pop III formation and Pop II formation with a normal IMF.

If correct, the implications of an ionizing background that is composed nearly equally of photons originating from Pop II and Pop III stars are interesting in several respects. The halos hosting the Pop II galaxies are rare peaks and will be highly clustered, while the lower-mass halos hosting the Pop III stars will be much less clustered, leading potentially to a rather complex topology for reionization. Because the Pop III stars will produce an even larger excess of helium ionizing photons, there are important implications for helium reionization as well (e.g. Venkatesan et al. 2003).

Acknowledgments

We thank L. Moustakas for a careful reading of the manuscript and useful suggestions.

REFERENCES

- Abel, T., Bryan, G. L., & Norman, M. L. 2000, *ApJ*, 540, 39
- . 2002, *Science*, 295, 93
- Barger, A. J., Cowie, L. L., & Richards, E. A. 2000, *AJ*, 119, 2092
- Becker, R. H. et al. 2001, *AJ*, 122, 2850
- Benson, A. J., Lacey, C. G., Baugh, C. M., Cole, S., & Frenk, C. S. 2002, *MNRAS*, 333, 156
- Bromm, V., Coppi, P. S., & Larson, R. B. 2002, *ApJ*, 564, 23
- Bromm, V., Kudritzki, R. P., & Loeb, A. 2001, *ApJ*, 552, 464
- Cen, R. 2002, preprint, astro-ph/0210473
- Ciardi, B., Ferrara, A., & White, S. D. M. 2003, preprint, astro-ph/0302451
- Cole, S., Aragón-Salamanca, A., Frenk, C., Navarro, J., & Zepf, S. 1994, *MNRAS*, 271, 781
- Cole, S., Lacey, C. G., Baugh, C. M., & Frenk, C. S. 2000, *MNRAS*, 319, 168
- Dekel, A. & Silk, J. 1986, *ApJ*, 303, 39

- Fan, X. 2001, *AJ*, 122, 2833
- Gnedin, N. Y. 2000, *ApJ*, 542, 535
- Haiman, Z. & Holder, G. 2003, preprint, astro-ph/0302403
- Hurwitz, M., Jelinsky, P., & Dixon, W. V. D. 1997, *ApJ*, 481, L31+
- Iwata, I., Ohta, K., Tamura, N., Ando, M., Wada, S., Watanabe, C., Akiyama, M., & Aoki, K. 2003, preprint, astro-ph/0301084
- Kauffmann, G., Colberg, J., Diaferio, A., & White, S. D. M. 1998, *MNRAS*, 303, 188
- Kauffmann, G., White, S., & Guiderdoni, B. 1993, *MNRAS*, 264, 201
- Kennicutt, R. 1983, *ApJ*, 272, 54
- . 1998, *ApJ*, 498, 181
- Kogut, A. et al. 2003, preprint, astro-ph/0302213
- Lanzetta, K. M., Yahata, N., Pascarelle, S., Chen, H., & Fernández-Soto, A. 2002, *ApJ*, 570, 492
- Larson, R. B. 1998, *MNRAS*, 301, 569
- Leitherer, C., Ferguson, H. C., Heckman, T. M., & Lowenthal, J. D. 1995, *ApJ*, 454, L19
- Leitherer, C. et al. 1999, *ApJS*, 123, 3
- Loeb, A. & Barkana, R. 2001, *ARA&A*, 39, 19
- Mac Low, M. & Ferrara, A. 1999, *ApJ*, 513, 142
- Machacek, M. E., Bryan, G. L., & Abel, T. 2001, *ApJ*, 548, 509
- . 2003, *MNRAS*, 338, 273
- Martin, C. L. 1999, *ApJ*, 513, 156
- Press, W. H. & Schechter, P. 1974, *ApJ*, 187, 425
- Ricotti, M., Gnedin, N. Y., & Shull, J. M. 2001, *ApJ*, 560, 580
- . 2002, *ApJ*, 575, 49
- Schaerer, D. 2003, *A&A*, 397, 527

- Somerville, R. S. 2002, *ApJ*, 572, L23
- Somerville, R. S. & Primack, J. R. 1999, *MNRAS*, 310, 1087
- Somerville, R. S., Primack, J. R., & Faber, S. M. 2001, *MNRAS*, 320, 504
- Spergel, D.N. et al. 2003, preprint, astro-ph/0302209
- Springel, V. & Hernquist, L. 2003, *MNRAS*, 339, 312
- Stanway, E., Bunker, A., & McMahon, R. 2003, preprint, astro-ph/0302212
- Steidel, C. C., Adelberger, K. L., Giavalisco, M., Dickinson, M., & Pettini, M. 1999, *ApJ*, 519, 1
- Steidel, C. C., Pettini, M., & Adelberger, K. L. 2001, *ApJ*, 546, 665
- Tegmark, M., Silk, J., Rees, M. J., Blanchard, A., Abel, T., & Palla, F. 1997, *ApJ*, 474, 1
- Thoul, A. A. & Weinberg, D. H. 1996, *ApJ*, 465, 608
- Tumlinson, J., Shull, J. M., & Venkatesan, A. 2003, *ApJ*, 584, 608
- Venkatesan, A., Tumlinson, J., & Shull, J. M. 2003, *ApJ*, 584, 621
- Wyithe, S. & Loeb, A. 2002, preprint, astro-ph/0209056
- . 2003, preprint, astro-ph/0302297
- Yoshida, N., Abel, T., Hernquist, L., & Sugiyama, N. 2003, preprint, astro-ph/0301645

Table 1: Summary of merger tree models

Model	τ_*	SN feedback	gas ejection	squelching	$z_{\text{reion}}^{(1)}$	z_{rec}	$z_{\text{reion}}^{(2)}$
MT-1a	constant	no	no	no	N/A	N/A	N/A
MT-1b	$\propto t_{\text{dyn}}$	no	no	no	N/A	N/A	N/A
MT-2a	$\propto t_{\text{dyn}}$	no	no	yes	20	N/A	N/A
MT-2b	$\propto t_{\text{dyn}}$	no	no	yes	15	N/A	N/A
MT-2c	$\propto t_{\text{dyn}}$	no	no	yes	15.5	13.5	6.0
MT-3a	$\propto t_{\text{dyn}}$	yes	no	yes	15.5	13.5	6.0
MT-3b	$\propto t_{\text{dyn}}$	yes	yes	yes	15.5	13.5	6.0
MT-3c	$\propto t_{\text{dyn}}$	yes	$V_c < 100 \text{ km/s}$	yes	15.5	13.5	6.0




Fig. 1.— The global star formation rate density as a function of redshift. Solid squares show observational estimates from direct integration of the rest-UV luminosity function of Lyman break galaxies at $z \sim 3$ and $z \sim 4$ from Steidel et al. (1999). The star symbol shows the estimate from the 850μ luminosity of sub-mm selected galaxies (Barger et al. 2000). The arrows at $z \sim 5$ and $z \sim 6$ show results from the detections of Lyman break galaxies from Iwata et al. (2003) and Stanway et al. (2003), respectively, where the bottom line shows the actual value detected and the top of the arrow shows the result of the (highly uncertain) correction for dust extinction and incompleteness (performed as described in SPF). Short, bold slanted lines show the three different estimates from Lanzetta et al. (2002), effectively reflecting different corrections for incompleteness due to cosmological surface brightness dimming. The SFRD predicted by the ‘halo-collapse’ models and the ‘merger tree models’ are also shown (see figure key and Table 1). In the left panel (a), the merger tree models do not include supernova feedback, and show different reionization histories. In the right panel (b), SN feedback has been included, and the treatment of gas ejection from the dark matter halos has been varied. The curve ending at $z \sim 20$ on the right panel shows the SFRD from numerical hydrodynamic simulations (Springel & Hernquist 2003).




Fig. 2.— The cumulative number of hydrogen ionizing photons per hydrogen atom in the Universe, produced by Pop II and Pop III stars. Bold solid and short-dashed lines show our ‘fiducial’ merger-tree models MT-3c and MT-3b, respectively, and the long dashed line shows the contribution from Pop III stars. The rate of production of ionizing photons by Pop II stars in galaxies overtakes that of Pop III stars sometime between $z \sim 20$ – 15 , perhaps triggering the reionization of the Universe.